

Understanding Upper Endoscopy

What is upper endoscopy?

Upper endoscopy lets your doctor examine the lining of the upper part of your gastrointestinal tract, which includes the esophagus, stomach and duodenum (first portion of the small intestine). Your doctor will use a thin, flexible tube called an endoscope, which has its own lens and light source, and will view the images on a video monitor. You might hear your doctor or other medical staff refer to upper endoscopy as upper GI endoscopy or esophagogastroduodenoscopy (EGD).

Why is upper endoscopy done?

Upper endoscopy helps your doctor evaluate symptoms of persistent upper abdominal pain, nausea, vomiting or difficulty swallowing. It's the best test for finding the cause of bleeding from the upper gastrointestinal tract. It's also more accurate than X-ray films for detecting inflammation, ulcers and tumors of the esophagus, stomach and duodenum.

Upper endoscopy is also used to treat conditions of the upper gastrointestinal tract. Your doctor can pass instruments through the endoscope to directly treat many abnormalities - this will cause you little or no discomfort. For example, your doctor might stretch (dilate) a narrowed area, remove polyps (usually benign growths) or treat bleeding.

What preparations are required?

An empty stomach allows for the best and safest examination. Your doctor will tell you when to start fasting as the timing can vary.

Tell your doctor in advance about any medications you take; you might need to adjust your usual dose for the examination. Discuss any allergies to medications as well as medical conditions, such as heart or lung disease.

Can I take my current medications?

Most medications can be continued as usual, but some medications can interfere with the preparation or the examination. Inform your doctor about medications you're taking, particularly aspirin products or antiplatelet agents, arthritis medications, anticoagulants (blood thinners such as warfarin or heparin), clopidogrel, insulin or iron products. Also, be sure to mention any allergies you have to medications.

What happens during upper endoscopy?

Your doctor might start by spraying your throat with a local anesthetic or by giving you a sedative to help you relax. You'll then lie on your side, and your doctor will pass the endoscope through your mouth and into the esophagus, stomach and duodenum. The endoscope doesn't interfere with your breathing.

What happens after upper endoscopy?

You will be monitored until most of the effects of the medication have worn off. Your throat might be a little sore, and you might feel bloated because of the air introduced into your stomach during the test. You will be able to eat after you leave unless your doctor instructs you otherwise.

If you have been given sedatives during the procedure, someone must drive you home and stay with you. Even if you feel alert after the procedure, your judgement and reflexes could be impaired for the rest of the day.

What are the possible complications of upper endoscopy?

Although complications can occur, they are rare when doctors who are specially trained and experienced in this procedure perform the test. Bleeding can occur at a biopsy site or where a polyp was removed, but it's usually minimal and rarely requires follow-up. Perforation (a hole or tear in the gastrointestinal tract lining) may require surgery but this is a very uncommon complication. Some patients might have a reaction to the sedatives or complications from heart or lung disease.

Although complications after upper endoscopy are very uncommon, it's important to recognize early signs of possible complications. Contact your doctor immediately if you have a fever after the test or if you notice trouble swallowing or increasing throat, chest or abdominal pain, or bleeding, including black stools. Note that bleeding can occur several days after the procedure.

If you have any concerns about a possible complication, it is always best to contact your doctor right away.

The above information was obtained from the ASGE website (<https://www.asge.org/home/for-patients/patient-information/understanding-upper-endoscopy>)